

Large/Massive Rotator Cuff Repair

Phase I (1-5 days post op)

- Wound care: Remove post op dressing at first P.T. appointment and replace with 4x4 gauze and paper tape. Monitor for signs of infection.
- Modalities: PRN for pain and inflammation.
- ROM:
 - o Initiate PROM of the shoulder as follows:
 - No shoulder adduction less than 30 degrees.
 - No shoulder flexion/abduction greater than 90 degrees for the first 6 weeks post op.
 - ER as tolerated.
 - No IR.
- Edema: Treat with modalities as appropriate.
- Sling: Sling worn continuously except during therapy or HEP sessions.
- Strengthening: Isometric grip strengthening.

Phase II (5 days-4 weeks post op)

- Wound care: Monitor for signs of infection.
- Modalities: PRN for pain and inflammation.
- ROM:
 - o Continue with Phase I
 - o Begin gentle IR in the scapular plane at 2 weeks post op.
- Edema: Continue to treat with modalities as appropriate.
- Sling: Continue sling use, including with abduction pillow, except for during therapy or HEP sessions until 6 weeks post op.
- Strengthening: Isometric grip strengthening only.

Phase III (4-10 weeks post op)

- ROM: at 6 weeks as follows:
 - o Initiate pulley exercises
 - o Initiate AROM
 - Achieve 120-140 degrees of flexion/abduction by 8 weeks post op.
 - Begin with supine, progress to reclined, followed by standing AROM for shoulder flexion.
 - Increase IR/ER as tolerated.
- Sling: Discontinue at 6 weeks post op.
- Strengthening: at 6 weeks post op as follows:
 - o Prone scapular stabilization.
 - o At the 5th week post op, may initiate sub-maximal, pain-free shoulder isometrics in all planes.
- Mobs: Glenohumeral joint mobilizations up to Grade III only.

Continued



Geoffrey Van Thiel, MD
www.vanthielmd.com

Phase IV (10 weeks + post op)

- ROM: Unrestricted
- Strengthening: Initiate PRE's
- Mobs: Glenohumeral joint mobilizations

Adapted From:

- 1.) Brotzman SB, Wilk KE. Clinical Orthopedic Rehabilitation Second Edition. Philadelphia: Mosby; 2002.
- 2.) Wilk, KE, Reinold MM, Andrews, JR. Type Three Rotator Cuff Repair Arthroscopic Assisted-Mini-Open Repair Large to Massive Tears (Greater than 4 cm). Winchester MA: Advanced Continuing Education Institute, 2004.