

# **Distal Triceps Repair (Advanced)**

Dr. Foster

Please follow the **Indiana Hand Protocol (5th edition)** for more conservative treatment. Therapy order will specify which post-op protocol to follow.

## Phase I (1 day - 6 week post-op)

- Begin therapy 2 weeks post-op
- Wound care: Initiate scar management as needed once incision site has healed
- Modalities: Initiate modalities as needed once incision site has healed
- Brace: Hinged elbow brace locked at 30 degrees of flexion
- ROM:
  - Gravity assisted AAROM elbow extension
  - Gentle AROM elbow flexion up to 90 degrees
    - Patient will perform these exercises within the hinged elbow brace. Clinician will educate patient on how to adjust/manipulate the brace.

### Phase II (6 weeks – 8 weeks post-op)

- Brace: Patient may begin weaning out of brace in controlled environments
- ROM:
  - Begin PROM for elbow flexion/extension
  - Full ROM allowed

### Phase III (8 weeks post-op)

- Brace: Discontinue hinged elbow brace
- Strengthening: May begin with MD clearance

### **Adapted From:**

1) Consultation with Dr. Brian Foster, MD at Ortholllinois

Initiation Date: 03/08/2023 Revised Date: N/A

324 Roxbury Road \* Rockford, IL \* Phone (815) 484-6990 \* Fax (815) 484-6961



Initiation Date: 03/08/2023 Revised Date: N/A

324 Roxbury Road \* Rockford, IL \* Phone (815) 484-6990 \* Fax (815) 484-6961